

MATHEMATICS

HOMOTOPY THEORY OF PRODUCTS ON SPHERES. I.

BY

P. W. H. LEMMENS

(Communicated by Prof. H. FREUDENTHAL at the meeting of January 25, 1969)

INTRODUCTION

In this thesis we study some special products on spheres from the standpoint of homotopy theory.

§§ 6–9 contain our main results. A survey of some results on reflecting products is found in theorem (9.21).

The map Gf , as defined in (4.2), is commonly known under the name “Hopf construction”, although this name is rather meaningless. Therefore we propose to call it “Hopf suspension”.

When considering S^n , we shall always exclude the case $n=0$.

By Z we denote the group of integers, and by Z_m we denote the group of integers modulo m .

The symbol \cong denotes that two groups are isomorphic.

The end of a proof is denoted by Q.E.D.

Numbers between square brackets refer to the references.

SP 1.2.3 stands for: E. H. Spanier, “Algebraic Topology”, chapter 1, section 2, item 3.

1. PRELIMINARIES

By (x_1, x_2, \dots, x_r) we denote the point $(y_1, y_2, \dots, y_k, \dots)$ in Hilbert space R , for which $y_k = x_k$ if $1 \leq k \leq r$ and $y_k = 0$ if $k > r$.

Let S^r be the subspace of R , which consists of the points $(x_1, x_2, \dots, x_{r+1})$ such that $x_1^2 + x_2^2 + \dots + x_{r+1}^2 = 1$.

$E_+^r, E_-^r \subset S^r$ are the hemispheres in which $x_{r+1} \geq 0, x_{r+1} \leq 0$.

The r -dimensional cube $I^r \subset R$ consists of the points (x_1, x_2, \dots, x_r) such that $-1 \leq x_i \leq 1, i = 1, 2, \dots, r$.

By \dot{I}^r we denote the boundary of I^r .

In all these spaces the basepoint will be the point

$$e = (-1, 0, \dots, 0, \dots).$$

We shall orient I^r ($r \geq 1$) as follows.

Let the points $a^{(i)}, i = 0, 1, 2, \dots, r$, be given by

$$a^{(i)} = (a_1^{(i)}, a_2^{(i)}, \dots, a_r^{(i)}).$$

Then the simplex $[a^{\langle 0 \rangle}, a^{\langle 1 \rangle}, \dots, a^{\langle r \rangle}]$ is positively oriented if and only if $(\det \{a_j^{\langle i \rangle} - a_j^{\langle 0 \rangle}; 1 \leq i \leq r, 1 \leq j \leq r\}) > 0$.

We choose the orientation of \dot{I}^r so, that \dot{I}^r is the combinatorial boundary of I^r .

Let the map $\varrho: S^{r-1} \rightarrow \dot{I}^r$ be the radial projection.

The orientation of S^{r-1} is such that the map ϱ has degree $+1$.

Let I denote the interval $-1 \leq t \leq 1$.

We shall often need a map

$$d_r: S^r \times I \rightarrow S^{r+1}$$

which has the following properties:

- (i) d_r maps $S^r \times (I - \dot{I})$ homeomorphically onto $S^{r+1} - e$,
- (ii) $d_r(S^r \times \dot{I} \cup e \times I) = e$,
- (iii) $d_r(x, 0) = x$ for all $x \in S^r$,
- (iv) $d_r(x, t) \in E_+^{r+1}$ if $t \geq 0$ and $d_r(x, t) \in E_-^{r+1}$ if $t \leq 0$, for all $x \in S^r$, $t \in I$.

Another important map is

$$\Phi_r: (I^r, \dot{I}^r) \rightarrow (S^r, e)$$

which is defined inductively by

$$\Phi_1(x) = (\cos \pi x, \sin \pi x), \quad x \in I^1,$$

$$\Phi_{n+1}(x_1, \dots, x_n, x_{n+1}) = d_n(\Phi_n(x_1, \dots, x_n), x_{n+1}).$$

Notice that Φ_r maps $I^r - \dot{I}^r$ homeomorphically onto $S^r - e$, and that Φ_r has degree $+1$.

It is very useful to represent elements of $\pi_r(X)$, where X is a space, in different ways. Usually one represents them by maps

$$(1.1) \quad \begin{aligned} f: (I^r, \dot{I}^r) &\rightarrow (X, x_0) \\ g: (S^r, e) &\rightarrow (X, x_0) \\ h: (\dot{I}^{r+1}, e) &\rightarrow (X, x_0) \end{aligned}$$

where x_0 is the basepoint of the space X .

We shall connect these representations by the following convention:

- (1.2) Let ϱ be the radial projection from S^r onto \dot{I}^{r+1} , and let f, g, h be maps as in (1.1).

If $f = g\Phi_r$ and $g = h\varrho$, then f, g, h represent the same element of $\pi_r(X)$.

When we consider a map or a homotopy, from a space X into a space Y , then we always assume that basepoints are respected.

Let X_1, X_2 and Y_1, Y_2 be subspaces of X and Y .

Then by the notations

$$(1.3) \quad \begin{aligned} f: (X, X_1, X_2) &\rightarrow (Y, Y_1, Y_2) \\ g: (X, X_1) &\rightarrow (Y, Y_1) \end{aligned}$$

we do not only mean that $f(X_1) \subset Y_1$, $f(X_2) \subset Y_2$, $g(X_1) \subset Y_1$, but also that these inclusions stay valid during homotopies of f and g .

2. THE FREUDENTHAL SUSPENSION

Consider a map $f: S^n \rightarrow S^r$.

Then there exists a map

$$Ef: S^{n+1} \rightarrow S^{r+1}$$

which has the following properties:

$$(2.1) \quad \begin{aligned} (i) \quad &Ef \text{ and } f \text{ agree on } S^n, \\ (ii) \quad &Ef(E_+^{n+1}) \subset E_+^{r+1} \text{ and } Ef(E_-^{n+1}) \subset E_-^{r+1}. \end{aligned}$$

It is well-known that, given f , all maps $S^{n+1} \rightarrow S^{r+1}$ which fulfil conditions (i) and (ii) of (2.1) are homotopic.

The *suspension homomorphism*

$$E: \pi_n(S^r) \rightarrow \pi_{n+1}(S^{r+1})$$

is now defined as follows: Let $f: S^n \rightarrow S^r$ represent $\alpha \in \pi_n(S^r)$, then $Ef: S^{n+1} \rightarrow S^{r+1}$ will be a representative map for the element $E\alpha \in \pi_{n+1}(S^{r+1})$.

By $[\alpha, \beta]$ we denote the Whitehead product between elements $\alpha \in \pi_p(X)$, $\beta \in \pi_q(X)$. For the definition of this product see § 3 of [19], where it is denoted by $\alpha \cdot \beta$.

Let ι_n be the positive generator of $\pi_n(S^n)$.

In [5] and [6] there occurs the sequence

$$(2.2) \quad \pi_i(S^n) \xrightarrow{E} \pi_{i+1}(S^{n+1}) \xrightarrow{H'} \pi_{i-n}(S^n) \xrightarrow{P} \pi_{i-1}(S^n) \xrightarrow{E} \dots$$

which is exact if $i \leq 3n-2$. In this sequence H' is a generalized Hopf invariant homomorphism, and P is the homomorphism defined by: $P\alpha = [\alpha, \iota_n]$, $\alpha \in \pi_{i-n}(S^n)$.

By Corollary (4.5) and Theorem (4.24) of [6], Corollary 2 of [5], and by the second corollary on page 247 of [17], we have for $k \geq 1$:

$$(2.3) \quad \begin{aligned} E: \pi_{2n}(S^n) &\rightarrow \pi_{2n+1}(S^{n+1}) \text{ is injective if } n=2 \text{ or } n=4k-1, \\ E: \pi_{2n-1}(S^{n-1}) &\rightarrow \pi_{2n}(S^n) \text{ is onto if } n=4k+2 \text{ or } n=4k+1. \end{aligned}$$

By the exactness of (2.2) and by the fact that $\pi_{n+1}(S^n) \cong Z_2$ for $n \geq 3$ (see [3]), we have in addition to (2.3):

$$(2.4) \quad \begin{aligned} \text{If } n=4k, \text{ then } E: \pi_{2n}(S^n) &\rightarrow \pi_{2n+1}(S^{n+1}) \text{ is not injective and} \\ E: \pi_{2n-1}(S^{n-1}) &\rightarrow \pi_{2n}(S^n) \text{ is not onto.} \end{aligned}$$

(2.5) **Lemma**

Let ι_n be the positive generator of $\pi_n(S^n)$, and let α be any element of $\pi_{2n}(S^n)$.

If n is *not* a multiple of 4, we have in $\pi_{2n}(S^n)$:

$$(k\iota_n) \circ \alpha = k\alpha \text{ for every integer } k.$$

PROOF: If n is not a multiple of 4, then it follows from (2.3) that either α is a suspension element, or $E: \pi_{2n}(S^n) \rightarrow \pi_{2n+1}(S^{n+1})$ is injective. (The case $n=1$ is fully trivial). If α is a suspension, then

$$(k\iota_n) \circ \alpha = k(\iota_n \circ \alpha) = k\alpha.$$

If E is injective, we form

$$E((k\iota_n) \circ \alpha - k\alpha) = k\iota_{n+1} \circ E\alpha - kE\alpha = kE\alpha - kE\alpha = 0.$$

Q.E.D.

3. SEPARATION ELEMENTS

The relative homotopy group $\pi_r(X, A)$, where A is a subspace of X , is defined to be the set of homotopy classes of maps

$$(I^r, \dot{I}^r, J^{r-1}) \rightarrow (X, A, x_0),$$

where $x_0 \in A$ is the basepoint of X , and J^{r-1} is the subspace of \dot{I}^r in which $x_r > -1$.

Every map $g: (I^r, \dot{I}^r, e) \rightarrow (X, A, x_0)$ is homotopic to a map

$$g': (I^r, \dot{I}^r, J^{r-1}) \rightarrow (X, A, x_0).$$

Now let Σ be the subspace of $S^p \times S^q$, defined by

$$\Sigma = S^p \times e \cup e \times S^q.$$

We orient $S^p \times S^q$ so, that the map

$$\Phi_{p,q}: (I^{p+q}, \dot{I}^{p+q}) \rightarrow (S^p \times S^q, \Sigma)$$

has degree $+1$. Here $\Phi_{p,q}$ is defined by

$$(3.1) \quad \Phi_{p,q}(x_1, \dots, x_p, x_{p+1}, \dots, x_{p+q}) = (\Phi_p(x_1, \dots, x_p), \Phi_q(x_{p+1}, \dots, x_{p+q})).$$

The positive generator γ of $\pi_{p+q}(S^p \times S^q, \Sigma)$ is then represented by a homotopic map

$$(3.2) \quad \Phi'_{p,q}: (I^{p+q}, \dot{I}^{p+q}, J^{p+q-1}) \rightarrow (S^p \times S^q, \Sigma, e \times e).$$

Now consider two maps $f, g: (S^p \times S^q, e \times e) \rightarrow (X, x_0)$ and assume that f and g are the same on Σ .

Then we define a map

$$D(f, g): (I^{p+q}, \dot{I}^{p+q}) \rightarrow (X, x_0)$$

by the formula

$$(3.3) \quad \begin{aligned} D(f, g)(x_1, \dots, x_{r-1}, x_r) &= f\Phi'_{p,q}(x_1, \dots, x_{r-1}, 2x_r - 1) \text{ if } x_r \geq 0, \\ &= g\Phi'_{p,q}(x_1, \dots, x_{r-1}, -2x_r - 1) \text{ if } x_r \leq 0, \end{aligned}$$

where $r = p + q$. The *separation element* $d(f, g) \in \pi_{p+q}(X)$ is now defined to be that element of $\pi_{p+q}(X)$ which is represented by $D(f, g)$.

For simplicity, let us assume that X is n -simple for all n (for example X arcwise connected and $\pi_1(X) = 0$), so the choice of the basepoint of X is immaterial.

Then, starting from (3.3), one easily sees that we can represent $d(f, g)$ also by a map

$$D'(f, g): S^{p+q} \rightarrow X$$

as follows. Take a map $u: (E_+^{p+q}, S^{p+q-1}) \rightarrow (I^{p+q}, \dot{I}^{p+q})$ of degree $+1$, and a map $v: (E_-^{p+q}, S^{p+q-1}) \rightarrow (I^{p+q}, \dot{I}^{p+q})$ of degree -1 , such that u and v agree on S^{p+q-1} .

Next we define $D'(f, g)$ by

$$(3.4) \quad \begin{aligned} D'(f, g)(x) &= f\Phi_{p,q}u(x) \text{ if } x \in E_+^{p+q}, \\ &= g\Phi_{p,q}v(x) \text{ if } x \in E_-^{p+q}. \end{aligned}$$

If we identify S^{p+q} and \dot{I}^{p+q+1} by the radial projection ϱ as in (1.2), then (3.4) would be a very useful formula if instead of u and v we could take the orthogonal projections along the x_{p+q+1} -axis from ϱE_+^{p+q} and from ϱE_-^{p+q} onto I^{p+q} .

However, these projections have degree $(-1)^{p+q}$ and $(-1)^{p+q+1}$ respectively.

For future purpose we define the map

$$D''(f, g): \dot{I}^{p+q+1} \rightarrow X$$

by the formula

$$(3.5) \quad \begin{aligned} D''(f, g)(x_1, \dots, x_r, x_{r+1}) &= f\Phi_{p,q}(x_1, \dots, x_r) \text{ if } x_{r+1} \geq 0, \\ &= g\Phi_{p,q}(x_1, \dots, x_r) \text{ if } x_{r+1} \leq 0, \end{aligned}$$

where $r = p + q$.

Because of our comment on (3.4), we have:

(3.6) **Lemma**

The map $D''(f, g): \dot{I}^{p+q+1} \rightarrow X$, as defined in (3.5), represents the element $(-1)^{p+q}d(f, g)$ of $\pi_{p+q}(X)$.

Now we supply a survey of the main properties of the separation element as these are listed by I. M. James in the appendix of [9], but we will write them down in the form we need later on. In this survey L stands for $S^p \times S^q$ and X, Y are spaces.

- (3.7) Let $f, g: L \rightarrow X$ be maps which agree on Σ . Then f is homotopic to g , relative to Σ , if and only if $d(f, g) = 0$.
- (3.8) Let $f, g, h: L \rightarrow X$ be maps which agree on Σ . Then $d(f, g) = d(f, h) + d(h, g)$.
- (3.9) Let an element $\delta \in \pi_{p+q}(X)$ and a map $f: L \rightarrow X$ be given. Then there exists a map $g: L \rightarrow X$, which agrees with f on Σ , such that $d(f, g) = \delta$.
- (3.10) Let $f_t, g_t: L \rightarrow X$ be homotopies which agree on Σ , where $0 \leq t \leq 1$. Then $d(f_0, g_0) = d(f_1, g_1)$.
- (3.11) Let $f, g: L \rightarrow X$ be maps which agree on Σ . For a map $h: X \rightarrow Y$, let $h_*: \pi_{p+q}(X) \rightarrow \pi_{p+q}(Y)$ be the homomorphism induced by h . Then $d(hf, hg) = h_*d(f, g)$.
- (3.12) Let $k: (L, \Sigma) \rightarrow (L, \Sigma)$ be a map of degree r . Let $f, g: L \rightarrow X$ be maps which agree on Σ . Then $d(fk, gk) = rd(f, g)$.

Property (3.12) calls for the following remark.

For the map $k: (L, \Sigma) \rightarrow (L, \Sigma)$, let $k'_*: H_{p+q}(L, \Sigma) \rightarrow H_{p+q}(L, \Sigma)$ be the induced homomorphism for integral relative homology groups. As the map $\Phi_{p,q}: (I^{p+q}, \dot{I}^{p+q}) \rightarrow (L, \Sigma)$ is a homeomorphism from $I^{p+q} - \dot{I}^{p+q}$ onto $L - \Sigma$, the integral relative homology groups $H_{p+q}(I^{p+q}, \dot{I}^{p+q})$ and $H_{p+q}(L, \Sigma)$ are isomorphic (see SP 4.8.9). Therefore we have:

$$H_m(L, \Sigma) \equiv H_m(I^m, \dot{I}^m) \equiv \mathbb{Z}, \text{ where } m = p + q.$$

Let α be a generator of $H_{p+q}(L, \Sigma)$, then the degree of the map k is said to be r , if $k'_*\alpha = r\alpha$.

In the exact homology sequence of the pair (L, Σ) applies:

$$j_*: H_{p+q}(L) \rightarrow H_{p+q}(L, \Sigma) \text{ is injective,}$$

since $H_{p+q}(\Sigma) = 0$.

It is well-known that $H_{p+q}(L) \equiv \mathbb{Z}$, with generator $\iota_p \otimes \iota_q$, the latter being the tensor product between generators ι_p and ι_q of $H_p(S^p)$ and $H_q(S^q)$ respectively.

Let $k''_*: H_{p+q}(L) \rightarrow H_{p+q}(L)$ be the homomorphism induced by the map k , and suppose that k has degree r .

Then we have: $k''_*(\iota_p \otimes \iota_q) = r(\iota_p \otimes \iota_q)$.

(3.13) **Theorem**

Let $u: S^p \rightarrow S^p$ and $v: S^q \rightarrow S^q$ be maps of degree m and n respectively. Then the map $k: (L, \Sigma) \rightarrow (L, \Sigma)$ defined by

$$k(x, y) = (u(x), v(y)) \quad x \in S^p, y \in S^q$$

has degree $m \cdot n$.

PROOF: $k_*(\iota_p \otimes \iota_q) = m\iota_p \otimes n\iota_q = m \cdot n(\iota_p \otimes \iota_q)$. Q.E.D.

We want to specialize property (3.11) in case X, Y are spheres:

(3.14) Let $f, g: L \rightarrow S^r$ be maps which agree on Σ , and suppose that the map $h: S^r \rightarrow S^n$ represents $\alpha \in \pi_r(S^n)$. Then $d(hf, hg) = \alpha \circ d(f, g)$.

In the appendix of [9], I. M. James defines the separation element up to a sign. Using our definition of the separation element, it is easily verified that the results of [9] are valid exactly, without changes of signs.

4. THE HOPF SUSPENSION

In the usual way we identify I^{m+n} with $I^m \times I^n$ and we consider $I^m \times \dot{I}^n$, $\dot{I}^m \times I^n$, $\dot{I}^m \times \dot{I}^n$ the corresponding subspaces of I^{m+n} .

We describe every point of \dot{I}^{p+q+2} by three co-ordinates (x, y, t) , $x \in S^p$, $y \in S^q$, $t \in I$, such that

(4.1) (x, y, t) denotes the point

$$((1-t) \cdot \varrho(x), \varrho(y)) \in I^{p+1} \times \dot{I}^{q+1} \text{ when } t \geq 0,$$

$$(\varrho(x), (1+t) \cdot \varrho(y)) \in \dot{I}^{p+1} \times I^{q+1} \text{ when } t \leq 0,$$

where ϱ is again the radial projection as in (1.2).

Notice that the correspondence (4.1) between $S^p \times S^q \times I$ and \dot{I}^{p+q+2} is homeomorphically on $S^p \times S^q \times (I - \dot{I})$.

For any map

$$f: S^p \times S^q \rightarrow S^r$$

we construct its *Hopf suspension*

$$Gf: \dot{I}^{p+q+2} \rightarrow S^{r+1}$$

by the formula

$$(4.2) \quad Gf(x, y, t) = d_r(f(x, y), t) \quad x \in S^p, y \in S^q, t \in I.$$

Let the map $\bar{f}: \dot{I}^{p+1} \times \dot{I}^{q+1} \rightarrow S^r$ be defined by

$$\bar{f}(\varrho(x), \varrho(y)) = f(x, y) \quad x \in S^p, y \in S^q.$$

Then we observe the following properties of the map Gf :

- (i) Gf agrees with \bar{f} on the subspace $\dot{I}^{p+1} \times \dot{I}^{q+1}$ of \dot{I}^{p+q+2} ,
- (ii) $Gf(\dot{I}^{p+1} \times \dot{I}^{q+1}) \subset E_+^{r+1}$ and $Gf(\dot{I}^{p+1} \times I^{q+1}) \subset E_-^{r+1}$.

Let $f: S^p \times S^q \rightarrow S^r$ be given, then any map $\tilde{f}^{p+q+2} \rightarrow S^{r+1}$ which fulfils properties (i) and (ii) above is homotopic to Gf .

Let $\{Gf\}$ denote the homotopy class of Gf in $\pi_{p+q+1}(S^{r+1})$.

Instead of $\{Gf\}$ we will often choose the element $c(f) \in \pi_{p+q+1}(S^{r+1})$, defined by

$$(4.3) \quad c(f) = (-1)^{q+1} \{Gf\},$$

according to Theorem (8.5) of [8].

Because of the last remark of § 3, we may copy Corollary (9.2) of [9] in the form we need:

(4.4) **Lemma**

Let $f, g: S^p \times S^q \rightarrow S^r$ be maps which agree on Σ . Then we have in $\pi_{p+q+1}(S^{r+1})$:

$$c(g) - c(f) = Ed(f, g).$$

A second important property of the Hopf suspension is

$$(4.5) \quad \text{If } f, g: S^p \times S^q \rightarrow S^r \text{ are homotopic maps, then } c(f) = c(g).$$

The type of a map $f: S^p \times S^q \rightarrow S^r$ is the pair (α, β) , where $\alpha \in \pi_p(S^r)$ and $\beta \in \pi_q(S^r)$ are the homotopy classes of the sections $f': S^p \rightarrow S^r$ and $f'': S^q \rightarrow S^r$, which are defined by $f'(x) = f(x, e)$ and $f''(y) = f(e, y)$ respectively.

$$(4.6) \quad (\text{see Theorem (2.24) of [8]})$$

Suppose we have maps $g: S^i \rightarrow S^p$ and $h: S^j \rightarrow S^q$.

Let $\alpha \in \pi_i(S^p)$ and $\beta \in \pi_j(S^q)$ denote the homotopy classes of g and h .

Let $f': S^i \times S^j \rightarrow S^r$ be the composition of a given map $f: S^p \times S^q \rightarrow S^r$ and the product map $g \times h: S^i \times S^j \rightarrow S^p \times S^q$.

Then $c(f') = c(f) \circ (\alpha \star \beta)$, where

$$\alpha \star \beta = (-1)^{p(j+q)} E^{p+1} \beta \circ E^{j+1} \alpha = (-1)^{i(j+q)} E^{q+1} \alpha \circ E^{i+1} \beta.$$

An easy consequence of (4.6) is

$$(4.7) \quad \text{If, for a map } f: S^p \times S^q \rightarrow S^r, f(x, y) \text{ depends only on } x, \text{ or only on } y, \text{ then } c(f) = 0.$$

$$(4.8) \quad \text{Let } f: S^p \times S^q \rightarrow S^r \text{ be a map, and suppose that the map } g: S^r \rightarrow S^n \text{ represents the element } \alpha \in \pi_r(S^n). \text{ Then we have in } \pi_{p+q+1}(S^{n+1}):$$

$$c(gf) = E\alpha \circ c(f).$$

Remark: (4.8) follows easily from (8.3) of [8], or from (4.2) and (4.3) above.

5. THE RIGHT SUSPENSION

In this section we follow the construction of H. Toda in § 3 of [13]. For a map $f: S^p \times S^q \rightarrow S^r$ there exists its *right suspension*

$$E'f: S^p \times S^{q+1} \rightarrow S^{r+1}$$

which has the following properties:

- (5.1) (i) $E'f$ agrees with f on the subspace $S^p \times S^q$ of $S^p \times S^{q+1}$,
(ii) $E'f(S^p \times E_+^{q+1}) \subset E_+^{r+1}$, $E'f(S^p \times E_-^{q+1}) \subset E_-^{r+1}$.

If the map $f: S^p \times S^q \rightarrow S^r$ is given, then all maps $S^p \times S^{q+1} \rightarrow S^{r+1}$ which fulfil conditions (i) and (ii) of (5.1) are homotopic.

In Lemma (3.2) of [13], Toda shows the connection between the right suspension and the Hopf suspension, by proving:

- (5.2) For any map $f: S^p \times S^q \rightarrow S^r$, we have in $\pi_{p+q+2}(S^{r+2})$:

$$\{G(E'f)\} = -E\{Gf\}.$$

According to (4.3) and (5.2) we can write down:

(5.3) **Lemma**

For any map $f: S^p \times S^q \rightarrow S^r$ we have in $\pi_{p+q+2}(S^{r+2})$:

$$c(E'f) = Ec(f).$$

In the following theorem we show the connection between the separation element and the right suspension.

(5.4) **Theorem**

Let $f, g: S^p \times S^q \rightarrow S^r$ be maps which agree on Σ , and let $E'f, E'g: S^p \times S^{q+1} \rightarrow S^{r+1}$ also have the same sections.

Then we have in $\pi_{p+q+1}(S^{r+1})$:

$$d(E'f, E'g) = Ed(f, g).$$

PROOF: According to (3.5) and lemma (3.6), $(-1)^{p+q+1}d(E'f, E'g)$ is represented by the map $F: \dot{I}^{p+q+2} \rightarrow S^{r+1}$, defined by

$$F(x_1, \dots, x_{p+q}, x_{p+q+1}, x_{p+q+2}) =$$

$$= E'f(\Phi_p(x_1, \dots, x_p), \Phi_{q+1}(x_{p+1}, \dots, x_{p+q+1})) \text{ if } x_{p+q+2} \geq 0,$$

$$= E'g(\Phi_p(x_1, \dots, x_p), \Phi_{q+1}(x_{p+1}, \dots, x_{p+q+1})) \text{ if } x_{p+q+2} \leq 0.$$

Now let $v: \dot{I}^{p+q+2} \rightarrow \dot{I}^{p+q+2}$ be the map which interchanges the last two co-ordinates, so

$$v(x_1, \dots, x_{p+q}, x_{p+q+1}, x_{p+q+2}) = (x_1, \dots, x_{p+q}, x_{p+q+2}, x_{p+q+1}).$$

The map v has degree -1 , thus $Fv: I^{p+q+2} \rightarrow S^{r+1}$ represents

$$(-1)^{p+q}d(E'f, E'g).$$

For clearness sake we write down Fv explicitly:

$$\begin{aligned} Fv(x_1, \dots, x_{p+q}, x_{p+q+1}, x_{p+q+2}) &= \\ &= E'f(\Phi_p(x_1, \dots, x_p), \Phi_{q+1}(x_{p+1}, \dots, x_{p+q}, x_{p+q+2})) \text{ if } x_{p+q+1} \geq 0, \\ &= E'g(\Phi_p(x_1, \dots, x_p), \Phi_{q+1}(x_{p+1}, \dots, x_{p+q}, x_{p+q+2})) \text{ if } x_{p+q+1} \leq 0. \end{aligned}$$

Now by the definitions of Φ_{q+1} , $E'f$, $E'g$ and $D''(f, g)$, we can sum up the following properties of Fv :

- (i) $Fv(x_1, \dots, x_{p+q+1}, 0) = D''(f, g)(x_1, \dots, x_{p+q+1})$,
- (ii) $Fv(x_1, \dots, x_{p+q+1}, x_{p+q+2}) \in E_+^{r+1}$ if $x_{p+q+2} \geq 0$,
- (iii) $Fv(x_1, \dots, x_{p+q+1}, x_{p+q+2}) \in E_-^{r+1}$ if $x_{p+q+2} \leq 0$.

So Fv satisfies the properties of $ED''(f, g)$ (see (2.1)), and from this we may conclude:

$$(-1)^{p+q}d(E'f, E'g) = (-1)^{p+q}Ed(f, g). \quad \text{Q.E.D.}$$

6. SOME GENERAL REMARKS ON PRODUCTS ON SPHERES

A product μ on S^n will be a map $\mu: S^n \times S^n \rightarrow S^n$, such that $\mu(e, e) = e$. For shortness sake we shall often note down: $\mu(x, y) = x \cdot y$.

Also in this section we shall use the notations:

$$\Sigma = S^n \vee S^n = S^n \times e \cup e \times S^n.$$

The product μ on S^n is said to be of type (p, q) , if the maps $\mu', \mu'': S^n \rightarrow S^n$, defined by $\mu'(x) = \mu(x, e)$ and $\mu''(x) = \mu(e, x)$, are of degree p, q respectively.

Let μ be a product on S^n , and let $f, g: X \rightarrow S^n$ be maps. Then we define the map $f \cdot g: X \rightarrow S^n$ by $f \cdot g(z) = f(z) \cdot g(z)$, $z \in X$.

(6.1) Theorem

Let μ be a product on S^n , and let $f, g: S^m \rightarrow S^n$ be maps. For a map h , $\{h\}$ denotes the homotopy class of h .

Then in $\pi_m(S^n)$ applies:

$$\{f \cdot g\} = \{f \cdot e\} + \{e \cdot g\},$$

where $e: S^m \rightarrow S^n$ denotes the constant map: $e(z) = e$.

PROOF: Represent $\{f\}, \{g\} \in \pi_m(S^n)$ by maps $f', g': (I^m, I^m) \rightarrow (S^n, e)$. Next consider the maps $F, G: (I^m, I^m) \rightarrow (S^n, e)$ defined by

$$\begin{aligned} F(x_1, x_2, \dots, x_m) &= e & \text{if } x_1 \geq 0, \\ &= f'(2x_1 + 1, x_2, \dots, x_m) & \text{if } x_1 \leq 0, \\ G(x_1, x_2, \dots, x_m) &= g'(2x_1 - 1, x_2, \dots, x_m) & \text{if } x_1 \geq 0, \\ &= e & \text{if } x_1 \leq 0. \end{aligned}$$

Now it is clear that $F \sim f'$ and $G \sim g'$, so $F \cdot G \sim f' \cdot g'$. Let us write down $F \cdot G: (I^m, \dot{I}^m) \rightarrow (S^n, e)$ explicitly:

$$\begin{aligned} F \cdot G(x_1, x_2, \dots, x_m) &= e \cdot g'(2x_1 - 1, x_2, \dots, x_m) \text{ if } x_1 \geq 0, \\ &= f' \cdot e(2x_1 + 1, x_2, \dots, x_m) \text{ if } x_1 \leq 0. \end{aligned}$$

Thus $\{F \cdot G\} = \{f' \cdot e\} + \{e \cdot g'\}$, and by the final remark, which is easy to prove, that $\{f \cdot g\}$ is independent of the representation of $\{f\}$ and $\{g\}$, the theorem is proved. Q.E.D.

For maps $f, g: S^n \times S^n \rightarrow S^n$, let $f|_\Sigma = g|_\Sigma$ denote that f and g agree on Σ .

(6.2) Theorem

Let μ be a product on S^n of type (p, q) , and let $u, v, u', v': (S^n \times S^n, e \times e) \rightarrow (S^n, e)$ be maps such that $u|_\Sigma = v|_\Sigma$ and $u'|_\Sigma = v'|_\Sigma$. Finally, let ι_n be the positive generator of $\pi_n(S^n)$.

Then in $\pi_{2n}(S^n)$ we have the equation:

$$d(u \cdot u', v \cdot v') = p\iota_n \circ d(u, v) + q\iota_n \circ d(u', v').$$

PROOF: Represent the separation element $d(u \cdot u', v \cdot v')$ by the map $D(u \cdot u', v \cdot v'): (I^{2n}, \dot{I}^{2n}) \rightarrow (S^n, e)$ as in (3.3). By simply writing down the explicit formula for $D(u \cdot u', v \cdot v')$ as in (3.3), it is immediately clear that

$$D(u \cdot u', v \cdot v') = D(u, v) \cdot D(u', v').$$

Now by theorem (6.1) we may note down:

$$\{D(u \cdot u', v \cdot v')\} = \{D(u, v) \cdot e\} + \{e \cdot D(u', v')\},$$

but $D(u, v) \cdot e = \mu' D(u, v)$ and $e \cdot D(u', v') = \mu'' D(u', v')$ (see the introduction of this section). By the fact that μ' and μ'' are maps of degree p and q , we have in $\pi_{2n}(S^n)$:

$$\{D(u \cdot u', v \cdot v')\} = p\iota_n \circ \{D(u, v)\} + q\iota_n \circ \{D(u', v')\}. \quad \text{Q.E.D.}$$

Because of lemma (2.5) we are able to simplify the assertion of this theorem, if n is not a multiple of 4:

(6.3) Corollary

If n is *not* a multiple of 4, then the assertion in theorem (6.2) reduces to

$$d(u \cdot u', v \cdot v') = p d(u, v) + q d(u', v').$$

Remark: Because of (3.14) the equation in theorem (6.2) is equivalent to the equation

$$d(u \cdot u', v \cdot v') = d(u \cdot e, v \cdot e) + d(e \cdot u', e \cdot v').$$

7. HOMOTOPY CLASSIFICATION OF PRODUCTS ON SPHERES

By the homotopy extension theorem, every map $f: S_1^n \times S_2^n \rightarrow S^n$ of type (p, q) is homotopic to an extension of a fixed map $g_0: S_1^n \vee S_2^n \rightarrow S^n$, such that $g_0|_{S_1^n}, g_0|_{S_2^n}$ have degree p, q respectively.

Because of this fact we can apply the theory of W. D. Barcus and M. G. Barratt in [2], in particular § 5 of [2].

We provide a brief account of the methods used in [2].

Let L be a CW complex, and let K be a subcomplex of L such that $L = K \cup e^{r+1}$. Here e^{r+1} is a $(r+1)$ -cel, and the attaching map will be $g: S^r \rightarrow K$.

For simplicity we assume here that $L = S^n \times S^n$ and $K = \Sigma$, so $r = 2n - 1$.

For a space X let X^K denote the function space of maps $(K, k_0) \rightarrow (X, x_0)$, where k_0, x_0 are basepoints.

Elements of $\pi_1(X^K, u_0)$, $u_0 \in X^K$, may be represented as homotopy classes of maps $F: (K \times S^1, k_0 \times S^1) \rightarrow (X, x_0)$ such that $F(k, e) = u_0(k)$.

The attaching map $g: S^r \rightarrow K$ induces a homomorphism

$$g^*: \pi_1(X^K, u_0) \rightarrow \pi_1(X^{S^r}, u_0g).$$

For any map $v: S^r \rightarrow X$ there is defined a map

$$v^b: (S^r \times S^1, e \times S^1) \rightarrow (X, x_0)$$

by $v^b(x, y) = v(x)$.

Let the element $\{F\} \in \pi_1(X^{S^r}, v_0)$ be represented by the map

$$F: (S^r \times S^1, e \times S^1) \rightarrow (X, x_0),$$

then the isomorphism

$$v_0^{\natural}: \pi_1(X^{S^r}, v_0) \rightarrow \pi_{r+1}(X, x_0)$$

is defined by $v_0^{\natural}\{F\} = d(F, v_0^b)$.

Let $\alpha \in \pi_r(K)$ be the homotopy class of the attaching map g , then there is defined a homomorphism

$$\alpha_{u_0}: \pi_1(X^K, u_0) \rightarrow \pi_{r+1}(X, x_0)$$

by the formula: $\alpha_{u_0} = (u_0g)^{\natural}g^*$.

Next there is proved in [2]:

Let $f_0, f_1: L \rightarrow X$ be extensions of $u_0: K \rightarrow X$, then we have: $f_0 \sim f_1$ if and only if $d(f_0, f_1) \in \alpha_{u_0}\pi_1(X^K, u_0)$.

Therefore, if the map $u_0: K \rightarrow X$ can be extended to L , the homotopy classes of extensions of u_0 are in 1-1 correspondence with the elements of $\pi_{r+1}(X, x_0)/\alpha_{u_0}\pi_1(X^K, u_0)$.

In case $L = S_1^n \times S_2^n$, $K = S_1^n \vee S_2^n$, $n > 1$, it turns out that the subgroup

$\alpha_{u_0}\pi_1(X^K, u_0)$ of $\pi_{r+1}(X, x_0)$ consists of the sums of Whitehead products

$$[\omega, \eta] + [\nu, \zeta]$$

where $\eta, \zeta \in \pi_{n+1}(X)$ are arbitrarily chosen, and where $\omega, \nu \in \pi_n(X)$ are represented by the maps $u_0|_{S_1^n}, u_0|_{S_2^n}$ respectively.

Let us now return to the case $L = S_1^n \times S_2^n$, $K = S_1^n \vee S_2^n$, $X = S^n$. First we make the following remark:

- (7.1) Let ι_n, η_n be generators of $\pi_n(S^n)$, $\pi_{n+1}(S^n)$ respectively.
Then $2[\iota_n, \eta_n] = 0$ when $n \geq 3$, for then is $\pi_{n+1}(S^n) \cong Z_2$.

It is now very easy to deduce from (7.1) and from [2] the following lemma:

(7.2) **Lemma**

Let $n \geq 3$ and let there be maps $S^n \times S^n \rightarrow S^n$ of type (p, q) . Then:

- a. The set of homotopy classes of maps $S^n \times S^n \rightarrow S^n$ of type (p, q) is in 1-1 correspondence with the elements of $\pi_{2n}(S^n)/W$, where W contains only the elements $0, p[\iota_n, \eta_n]$ and $q[\iota_n, \eta_n]$.
- b. Let $f, g: S^n \times S^n \rightarrow S^n$ be maps of type (p, q) which agree on Σ . Then f is homotopic to g , if and only if $d(f, g) \in W$.

Consider the suspension homomorphism $E: \pi_{2n}(S^n) \rightarrow \pi_{2n+1}(S^{n+1})$.

- (7.3) If $n \geq 3$, then the kernel of E consists precisely of the elements 0 and $[\iota_n, \eta_n]$.

The assertion in (7.3) is easily proved by taking $i = 2n + 1$ in the exact sequence (2.2).

Because of (7.1) and (7.3) we have in lemma (7.2):

- (7.4) If p, q are both even, then $\pi_{2n}(S^n)/W$ is isomorphic to $\pi_{2n}(S^n)$.
If p, q are not both even, then $\pi_{2n}(S^n)/W$ is isomorphic to $E\pi_{2n}(S^n)$.

A. *Homotopy classification of maps $S^n \times S^n \rightarrow S^n$ of type (p, q) , where $n \geq 3$ and p, q are both even.*

In this case the subgroup W of $\pi_{2n}(S^n)$ contains only the element 0 .

Take an arbitrary but fixed map $F: S^n \times S^n \rightarrow S^n$ of type (p, q) . For every map $f: S^n \times S^n \rightarrow S^n$ of type (p, q) we define an element $d_F(f) \in \pi_{2n}(S^n)$ as follows. Let f' be a map such that $f' \sim f$ and $f'|_{\Sigma} = F|_{\Sigma}$. We put

$$d_F(f) = d(F, f').$$

It follows from lemma (7.2) that $d_F(f)$ does not depend on the choice of f' within the homotopy class of f .

Furthermore, with the help of (3.8) it is easy to verify that the following lemma is valid:

(7.5) Lemma

Let $n \geq 3$, and let p, q be both even.

a. Let $F: S^n \times S^n \rightarrow S^n$ be a fixed map of type (p, q) .

Then the homotopy classes of maps $S^n \times S^n \rightarrow S^n$ of type (p, q) are fully determined by d_F .

b. Let $f, g: S^n \times S^n \rightarrow S^n$ be maps of type (p, q) . By choosing $F = f$, we have:

f is homotopic to g , if and only if $d_f(g) = 0$.

B. Homotopy classification of maps $S^n \times S^n \rightarrow S^n$ of type (p, q) , where $n \geq 3$ and p, q are not both even.

In this case it follows from (7.3) that the subgroup W of $\pi_{2n}(S^n)$ in lemma (7.2) is just the kernel of the suspension homomorphism E .

Therefore we can rewrite lemma (7.2) (b):

Let $f, g: S^n \times S^n \rightarrow S^n$ be maps of type (p, q) which agree on Σ , then $f \sim g$, if and only if $Ed(f, g) = 0$.

We take again an arbitrary but fixed map $F: S^n \times S^n \rightarrow S^n$ of type (p, q) . For each map f of type (p, q) we define an element $c_F(f) \in E\pi_{2n}(S^n)$ by:

$$c_F(f) = c(f) - c(F).$$

It follows from (4.5) that $c_F(f)$ is independent of f within its homotopy class.

Let $f, h: S^n \times S^n \rightarrow S^n$ be such that $c_F(f) = c_F(h)$. Then $c(f) = c(h)$.

Next we choose a map f' such that $f' \sim f$ and $f'| \Sigma = h| \Sigma$.

From lemma (4.4) we obtain:

$$c(f) - c(h) = c(f') - c(h) = Ed(h, f') = 0.$$

Hence $f' \sim h$, and thus $f \sim h$.

Therefore we can write down:

(7.6) Lemma

Let $n \geq 3$, and let p, q be not both even.

a. Let $F: S^n \times S^n \rightarrow S^n$ be a fixed map of type (p, q) .

Then the homotopy classes of maps $S^n \times S^n \rightarrow S^n$ of type (p, q) are fully determined by c_F .

b. Let $f, g: S^n \times S^n \rightarrow S^n$ be maps of type (p, q) .

Then f is homotopic to g , if and only if $c(f) = c(g)$.

C. Homotopy classification of maps $S^n \times S^n \rightarrow S^n$ of type (p, q) , where $n = 2$ or $n = 4k - 1$ and p, q are arbitrary.

Let $F: S^n \times S^n \rightarrow S^n$ be a fixed map of type (p, q) . Let f, f' be maps of type (p, q) such that $f \sim f'$ and $f| \Sigma = f'| \Sigma = F| \Sigma$.

Then, by (4.4) and (4.5), we have:

$$Ed(F, f) = c(f) - c(F) = c(f') - c(F) = Ed(F, f').$$

So $d(F, f) = d(F, f')$, since E is injective by (2.3).

Therefore we can form $d_F(f)$ and $c_F(f)$ for any map $f: S^n \times S^n \rightarrow S^n$ of type (p, q) , and the relation between them is: $E d_F(f) = c_F(f)$.

Let f, g be maps of type (p, q) such that $f \sim g$, then $c_F(f) = c_F(g)$ by (4.5), so $d_F(f) = d_F(g)$.

For maps f, g let $c_F(f) = c_F(g)$, then $d_F(f) = d_F(g)$, so $f \sim g$ by (3.7).

(7.7) **Lemma**

Let $n=2$ or let $n=4k-1$.

a. Let $F: S^n \times S^n \rightarrow S^n$ be a fixed map of type (p, q) .

Then the homotopy classes of maps $S^n \times S^n \rightarrow S^n$ of type (p, q) are fully determined by c_F , as well as by d_F .

b. Let $f, g: S^n \times S^n \rightarrow S^n$ be maps of the same type.

Then the following relations are equivalent:

$$f \sim g, \quad d_f(g) = 0, \quad c(f) = c(g).$$

To complete this classification, we remark:

(7.8) Let $f, g: S^1 \times S^1 \rightarrow S^1$ be maps of the same type.

Then f is homotopic to g .

For choose a map $f': S^1 \times S^1 \rightarrow S^1$ such that $f' \sim f$ and $f'|_{\Sigma} = g|_{\Sigma}$. Then $d(f', g) = 0$, since $\pi_2(S^1) = 0$. Hence $f' \sim g$ and thus $f \sim g$.

(To be continued)